

RESEARCH ARTICLE

Effects of arbuscular mycorrhizal fungi, poultry manure compost, and cadmium on plant growth and nutrient absorption of *Oryza sativa*

Geomar Vallejos-Torres^{1,2*}, Nery Gaona-Jimenez¹, Luis Ordoñez-Sánchez¹, Carlos Lozano¹, Alberto Alva Arevalo³, Andi Lozano³, Wilfredo Mendoza-Caballero⁴, Christopher Paredes⁵, Jorge Saavedra-Ramírez⁵, James Tuanama⁶, and César Marín^{2,7}

¹Universidad César Vallejo, Catachachi, San Martín, Perú.

²Universidad Santo Tomás, Centro de Investigación e Innovación para el Cambio Climático (CiiCC), Valdivia, Chile.

³Universidad Nacional de San Martín, Jr. Maynas N°177, Tarapoto, San Martín, Perú.

⁴Universidad Católica Sedes Sapientiae, Los Olivos, Lima, Perú.

⁵Universidad Nacional Autónoma de Alto Amazonas (UNAAA), Alto Amazonas, Yurimaguas, Perú.

⁶Universidad Nacional Agraria La Selva, Tingo María, Huánuco, Perú.

⁷Vrije Universiteit Amsterdam, Amsterdam Institute for Life and Environment, Amsterdam, The Netherlands.

*Corresponding author (gvallejost@gmail.com).

Received: 4 April 2023; Accepted: 20 June 2023, doi:10.4067/S0718-58392023000600656

ABSTRACT

Compost produced with chicken manure and vegetable residues enriched with arbuscular mycorrhizal fungi (AMF) could become a sustainable agricultural strategy in biofertilization and bioremediation in rice (*Oryza sativa* L.) A completely randomized pot experiment was carried out to investigate the effects of AMF, humified composted poultry manure, and Cd on vegetative growth and elemental uptake of rice. After 90 d, rice treated with a combination of AMF and compost showed the highest plant growth. Overall, the best values in plant height, root length, chlorophyll content, and cation exchange capacity were obtained with the application of AMF and compost. In the absence of compost, Cd contamination significantly reduced AMF root colonization, while AMF alone increased root N, P, and K. Soil organic matter was higher when AMF, compost, and Cd were combined. Total soil P increased significantly with AMF inoculation (solely) compared to the rest of the treatments. The highest plant height (57.77 cm), root length (31.67 cm) and weight (167 g), shoot weight (74.67 g), and chlorophyll content (34.30 SPAD units) were achieved in the AMF-compost treatment. The highest values of root N (1.037%) and soil organic matter (2.66%) were obtained with the Compost-Cd treatment, which in turn presented lower Cd contents in roots and shoots. Cadmium inhibited morphological growth of the plant, biomass accumulation, and chlorophyll, but these parameters improved significantly with the application of AMF and compost, by facilitating the absorption of nutrients, particularly P.

Key words: Arbuscular mycorrhizal fungi, cadmium, *Oryza sativa*, synergistic effects.

INTRODUCTION

Rice (*Oryza sativa* L.) is a crop food of great nutritional value for mankind, being a critical crop in global food security. In Peru -and worldwide, rice is part of the essential family diet (Liu et al., 2014), and its cultivation is one of the main agricultural products: in the last 10 yr, the national production of paddy rice increased from 1 962 000 to 2 783 000 t, which represents an average annual increase rate of 4.2% (Neira et al., 2020). The Region of San Martín is considered the Peruvian region with the highest rice production, whose harvested area is 101 255 ha, with an average yield of 7 t ha⁻¹ (MINAGRI, 2017). This extensive area

planted with rice has demanded high levels of water use; likewise, agricultural soils contaminated by Cd, caused mainly by mining, electronic waste, and excessive use of N and P fertilizers, has been found in large-scale agricultural soils (Chen et al., 2019). Cadmium is absorbed by plants, accumulating in rice grains and entering human bodies due to its long half-life of up to 25 to 30 yr (Song et al., 2015), representing a serious threat to food security and human health (Li et al., 2019; Saikat et al., 2022). Therefore, soil contamination by heavy metals such as Cd not only inhibits plant growth, but also threatens human health, through food chain bioaccumulation (Zhang et al., 2019).

Cadmium content in rice soil has been reported up to 0.300-1.112 mg kg⁻¹ (Song et al., 2015), while in grains the values were up to 0.062 ± 0.128 mg kg⁻¹ (Song et al., 2017). Rice consumption has been considered the most important route of Cd exposure in humans (Li et al., 2017). Cadmium accumulation in rice is remarkably influenced by soil physical and chemical properties, such as soil organic matter (SOM), cation exchange capacity (CEC), pH, and other elements.

Faced with this problem, soil microbes play an essential role by affecting the characteristics of the soil and plant growth, which significantly influences the absorption of heavy metals. Soil microorganisms have been used over the last decades in order to increase plant growth, especially under soil limiting conditions. Spores of arbuscular mycorrhizal fungi (AMF) - beneficial soil microorganisms that form mutualistic symbioses with roots of most higher plant species, with the capacity to immobilize cadmium in the soil-plant system, have been used due to their strategies of immobilization and absorption of metals in the hyphae (Janeeshma and Puthur, 2020). They can increase root surface area for nutrient and water uptake (Liu et al., 2018). Likewise, compost prepared with manure has been considered as an important input to mitigate Cd toxicity in soils. Many studies show that composted manure immobilizes heavy metals in soil and reduces its uptake by plants; in turn, they improve soil conditions and plant growth, constituting a good alternative for heavy metal remediation. The main objective of this study was to elucidate the effects of AMF, poultry-manure compost, and Cd on plant growth and nutrient absorption in *Oryza sativa*.

MATERIALS AND METHODS

Soil and organic compost characteristics

The agricultural soil used (a total of 72 kg sieved soil) comes from fields close to large extensions of rice planted in the San Martín region, Peru, at a 0-20 cm soil depth. The chemical characteristics of the soil were: Total N, 1%; available (Olsen) P, 5.12 mg kg⁻¹; available K, 71.23 mg kg⁻¹; soil organic matter (SOM), 2.46%; total Cd, 0.31 mg kg⁻¹; pH, 7.18; and cation exchange capacity (CEC), 396.20 μS cm⁻¹. The humified compost was made with free-range hen manure and organic field residues and at the same time, it was enriched with earthworm humus and had the following properties: Total N, 2.12%; available P, 1300 mg kg⁻¹; available K, 15900 mg kg⁻¹; organic matter, 28.63%; total Cd, 0.00 mg kg⁻¹; pH, 7.12; and CEC, 1789.23 μS cm⁻¹. For the treatments contaminated with Cd, an aqueous solution of CdCl₂ (5 mg kg⁻¹) was added to each pot containing 3 kg substrate. To help stabilize the contaminant, the Cd-containing soil was allowed to air-dry for 50 d, and it was agitated every 15 d (four times), following the methodology of Vallejos-Torres et al. (2022).

Experimental design

This experiment consisted of a 2 × 2 × 2 factorial design with the addition of Cd (0 and 5 mg kg⁻¹), addition of humified compost ('Compost' thereafter) (0 and 250 g plant⁻¹, w:w), and inoculation with arbuscular mycorrhizal fungi (AMF) (control without mycorrhizal fungi and inoculation with AMF at 8.33%) arranged in a completely randomized design with three replicates. The AMF inoculum consisted of 2000 spores isolated from the Mariscal Cáceres Province in the San Martín region, Peru. The mycorrhizal inoculum was composed of a mixture of spores of the following species: *Claroideoglossum* sp., *Microkamienskia peruviana*, *Microkamienskia* sp., *Diversispora* sp., and *Claroideoglossum etunicatum* (Vallejos-Torres et al., 2022).

The growth substrate consisted of a mixture of river sand and agricultural soil (1:2, w:w), which was sterilized by autoclaving at 120 °C for 2 h. The seeds of rice 'INIA 507 - La Conquista' were disinfected with 2.5% sodium hypochlorite and distilled water; then, four seeds were sown per individual pot, filled

each with 3.0 kg growth substrate. The experiment was carried out in a greenhouse, located at the National University of San Martín in Tarapoto (Peru) (06°28'59.71" S, 76°21'18.90" W) from January to April 2022. During the experiment, the air temperature was 25-34 °C. The plants were irrigated regularly with sterile water at 60% of the field capacity, without any fertilization treatment.

Sampling and measurements

After 90 d, a total of 96 plants (eight treatments × three replicates × four plants per replicate) were harvested. Plant roots were washed and weighed on an analytical balance to determine fresh biomass and Cd by the HNO₃/spectroscopy digestion method with atomic absorption (Isaac and Johnson, 1975). Both roots and shoots were extracted from each plant sample to quantify Cd concentration (digestion with HNO₃) using atomic absorption spectroscopy (Zhang et al., 2020), and another group of shoots were taken to the laboratory for foliar analysis of N, using the Kjeldahl method (Kjeldahl 1883). Phosphorus was measured by the digestion of HNO₃/spectroscopy UV-Vis ($\lambda = 420$ nm) method, and K by the HNO₃/spectroscopy digestion method with atomic absorption (Isaac and Johnson, 1975). Substrate pH was measured in an aqueous extract (1:5) using a pH-meter; for the determination of SOM, the Walkley-Black method (De Vos et al., 2007) was followed. The CEC was determined by potential acidity + sum of bases, and the available P was estimated according to the Olsen method (Olsen and Sommers, 1982). Rhizospheric soil was also collected for the measurement of mycorrhizal root colonization. The height of the plant was measured from the base to the highest tip of the leaf. Arbuscular mycorrhizal colonization of roots was measured by the grid line intersection method after cleaning roots in 10% (w/v) KOH and staining them, as described by Phillips and Hayman (1970). Mycelium lengths per unit weight of soil were determined using Newman's formula (Newman, 1966).

Statistical analysis

All analyses were performed using the R programming language version 4.0.2 (R Core Team, R Foundation for Statistical Computing, Vienna, Austria). The normality and homoscedasticity of the data were verified using the Shapiro Wilk and Breush-Pagan tests, respectively. To analyze the effect of the treatments (AMF, compost, Cd, and their combinations) on the measured parameters, data were analyzed through ANOVA (using the R base function "aov") and Tukey-mean comparison tests with a probability of error of 5%, using the R base function "TukeyHSD".

RESULTS AND DISCUSSION

Effects on plant growth and chlorophyll

Root length and mycorrhizal colonization were significantly reduced when Cd was applied (Figure 1). As expected, root length was higher when both AMF and compost were combined (Figure 1a), while -and also as expected- AMF root colonization was higher when these fungi were applied alone (Figure 1b). Following the same pattern, a combination of AMF and compost resulted in significantly higher plant height (Figure 2a), shoot (Figure 2b) and root (Figure 2c) biomass, and chlorophyll content (Figure 2d). In all these cases, Cd applied alone resulted in trait values as low (or lower) as the control (Figure 2). The addition of compost and of compost + AMF to a Cd-contaminated pot, always resulted in significant increases in plant height, shoot and root biomass, and chlorophyll content (Figure 2). In most traits, except plant height, applying AMF alone resulted in increases compared to applying compost alone (Figure 2).

The interaction between AMF inoculum, poultry manure compost, and Cd had significant effects ($p < 0.001$) on plant height, root length, root biomass, and shoot biomass of *O. sativa*. Growth indicators and biomass characteristics also improved significantly with the application of compost made with chicken manure compared to the control. Chicken manure could provide essential nutrients directly to rice as indicated by Liu et al. (2021) when studying this waste material in maize, and its effects in plant growth. Without Cd contamination, root and shoot biomasses increased significantly. Cadmium contamination negatively affected root mycorrhizal colonization. This coincides with Wu et al. (2016), whom show negative effects of Cd (and Pb) toxicity on AMF and root endophytic fungi associated with wheat. Similar negative effects were shown by Vallejos-Torres et al. (2022) in cacao plants. Plant height and root length

were higher under AMF, compost, and AMF + compost treatments, but were negatively affected by Cd application, as roots are the main way in which Cd enters to the plant, hindering its growth (Kanu et al., 2017). Cadmium content in rice roots was quite higher than in shoots, coinciding with Kanu et al. (2017), although the same authors indicate that Cd transport in the aerial parts of the plant can vary between cultivars. Similarly, Li et al. (2019) demonstrated that Cd significantly inhibited root length and dry weight in rice seedlings. Cd toxicity minimizes the mitotic division of meristematic cells, leading to a reduction in root growth and length and dry biomass (Barman et al., 2020).




Figure 1. Effects of arbuscular mycorrhizal fungi (AMF) inoculation, humidified compost modification, Cd addition, and their combination on rice root length (a) and mycorrhizal colonization (b). Lowercase letters indicate significant differences between the different treatments.

The highest chlorophyll contents were found in the AMF and AMF + compost treatments; rice plants exposed to Cd showed inhibition of chlorophyll in their leaves (Herath et al., 2015). Accumulation of Cd in leaves could impair the chlorophyll biosynthesis and, therefore, reduce the chlorophyll content (Kanu et al., 2017). Furthermore, some authors such as Farooq et al. (2022) and Li et al. (2021) reported that Cd stress led to a substantial reduction in chlorophyll content in different rice varieties. The damage of chlorophyll pigments, photosynthesis, transpiration, stomatal conductance, and other physiological and photosynthetic processes, is due to deterioration in the thylakoid membrane and the formation of chlorophyll precursors (Vaculík et al., 2015).




Figure 2. Effects of arbuscular mycorrhizal fungi (AMF) inoculation, humidified compost modification, Cd addition, and their combination on rice plant height (a) shoot biomass (b), root biomass (c), and chlorophyll content (d). Lowercase letters indicate significant differences between the different treatments.

Effects on nutrient absorption

Interestingly, and in contrast with the marked effects and patterns found for AMF, compost, and Cd application on rice root length and mycorrhizal colonization (Figure 1) and plant height, root and shoot biomass, and chlorophyll (Figure 2), such patterns were not found for rice shoot nutrient absorption (Figure 3). Overall, there were nonsignificant differences regardless of treatment for rice shoot N (Figure 3a), P (Figure 3c), and K (Figure 3e). When compost was combined with AMF and, unexpectedly, with Cd, higher root N values were obtained (Figure 3b). The AMF alone and combined with compost resulted in higher root P and K (Figures 3d, 3f). The Cd significantly reduced the efficiency of compost and AMF in capturing root N, P, and K (Figure 3).




Figure 3. Effects of arbuscular mycorrhizal fungi (AMF) inoculation, humidified compost modification, Cd addition, and their combination on rice shoot N (a), root N (b), shoot P (c), root P (d), shoot K (e), and root K (e). Lowercase letters indicate significant differences between the different treatments.

The treatment combination of AMF and compost positively affected N, P, and K content in both rice roots and shoots. Previous studies have shown that AMF can facilitate the uptake of mineral elements by plants, particularly P (Liu et al., 2017). This is well established in mycorrhizal ecology, as hyphae explore a greater soil volume than roots by themselves (Liu et al., 2017). Root and shoot N, P, and K contents were higher than the control after applying compost, even in Cd-contaminated soils. Cadmium hindered the concentration of these nutrients in rice shoots and roots. The AMF inoculation in combination with the compost amendment had significant synergistic benefits, not only for nutrient uptake but also for the reduction in Cd uptake of rice in Cd-contaminated soils, results similar to those obtained by Zhang et al. (2019).

Effects on Cd and soil nutrients

All Cd-contaminated treatments had similar values of shoot Cd in rice plants (Figure 4a). Root Cd was the highest when Cd was applied alone, followed by when it was applied with AMF, then with AMF and compost, and finally, just with compost (Figure 3b).




Figure 4. Effects of arbuscular mycorrhizal fungi (AMF) inoculation, humidified compost modification, Cd addition, and their combination on rice shoot Cd (a) and root Cd (b). Lowercase letters indicate significant differences between the different treatments.

The results for the Tukey tests indicate that soil pH was lower in the control treatment; however, when AMF, compost, and Cd were added, they showed significant differences (Table 1). The SOM had similar results, showing the highest content when AMF + Cd + compost were combined, and the lowest SOM value was obtained by the control treatment. The highest CEC was presented by the treatment with the application of AMF and the application of compost, showing significant differences with the control treatment (lowest CEC value). Available P increased significantly by AMF inoculation compared to the control treatment; however, when Cd was added, it was reduced to a nonsignificant level.

Table 1. Tukey tests for soil chemical parameters across the different treatments. SOM: Soil organic matter; CEC: cation exchange capacity; AMF: arbuscular mycorrhizal fungi.

Treatment	pH	SOM %	CEC meq 100 g ⁻¹	Olsen P mg kg ⁻¹
Control	6.16 ± 0.21 ^c	1.29 ± 0.10 ^c	5.75 ± 0.03 ^c	4.95 ± 0.10 ^c
Compost	7.04 ± 0.08 ^b	1.85 ± 0.12 ^{bc}	8.15 ± 0.32 ^a	6.71 ± 0.15 ^b
AMF	7.44 ± 0.01 ^a	2.38 ± 0.15 ^{ab}	8.10 ± 0.06 ^a	7.33 ± 0.18 ^a
Cd	7.10 ± 0.05 ^b	1.92 ± 0.12 ^{bc}	7.70 ± 0.23 ^{ab}	6.84 ± 0.16 ^{ab}
Compost-Cd	7.43 ± 0.01 ^a	2.66 ± 0.12 ^a	7.70 ± 0.17 ^{ab}	7.24 ± 0.07 ^{ab}
Compost-AMF	7.57 ± 0.00 ^a	2.36 ± 0.14 ^{ab}	7.65 ± 0.26 ^{ab}	7.12 ± 0.00 ^{ab}
AMF-Cd	7.42 ± 0.02 ^a	1.42 ± 0.02 ^c	7.00 ± 0.00 ^b	6.89 ± 0.00 ^{ab}
Compost-Cd-AMF	7.49 ± 0.05 ^a	2.49 ± 0.02 ^a	8.00 ± 0.12 ^a	7.00 ± 0.07 ^{ab}

It is known that the addition of compost stabilizes soil pH through a liming effect that also favors plant growth (Ahmad et al., 2014). Compost addition significantly increased soil pH under AMF inoculation. Chicken manure is an organic material that has recently attracted attention for its higher levels of fertility (Wang et al., 2020). It is a stabilized soil amendment that can improve and restore SOM, improve water retention, and soil structure. In the same venue, CEC was higher under the AMF + compost treatment, and we think this treatment allows roots to growth by increasing pH, available P, and CEC, among other properties (Pandian et al., 2016). Likewise, mycorrhizal associations can contribute to reducing the transfer of heavy metals to plants by acting as an exclusion barrier (Cabral et al., 2015), which can react by binding heavy metals to the fungal hyphae (Gonzalez-Chavez et al., 2002).

CONCLUSIONS

Cadmium inhibited plant morphological growth and biomass accumulation in rice, probably caused by metal stress in rice roots. Inoculation with arbuscular mycorrhizal fungi (AMF) alone or with poultry manure compost significantly improved the rice growth. The AMF can facilitate plant nutrient uptake, particularly P. Likewise, Cd is known to inhibit plant nutrient uptake within the rhizosphere in rice. The results showed that Cd toxicity resulted in a substantial reduction of chlorophyll content while the addition of poultry manure compost significantly improved AMF colonization, and this in turn significantly increased root P concentration.

Author contributions

Conceptualization: G.V-T., N.G-J. Methodology: G.V-T., C.M. Software: G.V-T. Validation: C.M., L.A.O-S. Formal analysis: C.L., C.M. Investigation: A.A.A. Resources: A.L. Data curation: G.V-T., W.M-C., C.M. Writing-original draft: C.P. Writing-review & editing: J.S-R, C.M. Visualization: L.A., C.M. Supervision: N.G-J. Project administration: G.V-T. Funding acquisition: J.T. All co-authors reviewed the final version and approved the manuscript before submission.

Acknowledgements

G.V.T. thanks the agroforestry research center of Perú. C.M. thanks the Convocatoria Nacional Subvención a Instalación en la Academia Convocatoria Año 2021 + Folio SA77210019 (ANID-Chile).

References

- Ahmad, M., Lee, S.S., Lim, J.E., Lee, S.E., Cho, J.S., Moon, D.H., et al. 2014. Speciation and phytoavailability of lead and antimony in a small arms range soil amended with mussel shell, cow bone and biochar: EXAFS spectroscopy and chemical extractions. *Chemosphere* 95:433-441. doi:10.1016/j.chemosphere.2013.09.077.
- Barman, F., Majumdar, S., Arzoo, S.H., Kundu, R. 2020. Genotypic variation among 20 rice cultivars/landraces in response to cadmium stress grown locally in West Bengal, India. *Plant Physiology and Biochemistry* 148:193-206. doi:10.1016/j.plaphy.2020.01.019.

- Cabral, L., Soares, C.R., Giachini, A.J., Siqueira, J.O. 2015. Arbuscular mycorrhizal fungi in phytoremediation of contaminated areas by trace elements: mechanisms and major benefits of their applications. *World Journal of Microbiology and Biotechnology* 31:1655-1664. doi:10.1007/s11274-015-1918-y.
- Chen, J., Zou, W., Meng, L., Fan, X., Xu, G., and Ye, G. 2019. Advances in the uptake and transport mechanisms and QTLs mapping of cadmium in rice. *International Journal of Molecular Sciences* 20(14):3417. doi:10.3390/ijms20143417.
- De Vos, B., Lettens, S., Muys, B., Deckers, J.A. 2007. Walkley-Black analysis of forest soil organic carbon: recovery, limitations and uncertainty. *Soil Use and Management* 23:221-229. doi:10.1111/j.1475-2743.2007.00084.x.
- Farooq, M.U., Ishaq, I., Barutcular, C., Skalicky, M., Maqbool, R., Rastogi, A., et al. 2022. Mitigation effects of selenium on accumulation of cadmium and morpho-physiological properties in rice varieties. *Plant Physiology and Biochemistry* 170:1-13. doi:10.1016/j.plaphy.2021.11.035.
- Gonzalez-Chavez, C., D'Haen, J., Vangronsveld, J., Dodd, J.C. 2002. Copper sorption and accumulation by the extraradical mycelium of different *Glomus* spp. (arbuscular mycorrhizal fungi) isolated from the same polluted soil. *Plant and Soil* 240:287-297. doi:10.1023/A:1015794622592.
- Herath, I., Kumarathilaka, P., Navaratne, A., Rajakaruna, N., Vithanage, M. 2015. Immobilization and phytotoxicity reduction of heavy metals in serpentine soil using biochar. *Journal of Soils and Sediments* 15:126-138. doi:10.1007/s11368-014-0967-4.
- Isaac, R.A., Johnson, W.C. 1975. Collaborative study of wet and dry ashing techniques for the elemental analysis of plant tissue by atomic absorption spectrophotometry. *Journal of Association of Official Analytical Chemists* 58:436-440. doi:10.1093/jaoac/58.3.436.
- Janeeshma, E., Puthur, J.T. 2020. Direct and indirect influence of arbuscular mycorrhizae on enhancing metal tolerance of plants. *Archives of Microbiology* 202:1-16. doi:10.1007/s00203-019-01730-z.
- Kjeldahl, J. 1883. A new method for the determination of nitrogen in organic matter. *Zeitschrift für Analytische Chemie* 22:366-382. doi:10.1007/BF01338151.
- Kanu, A.S., Ashraf, U., Mo, Z., Fuseini, I., Mansaray, L.R., Duan, M., et al. 2017. Cadmium uptake and distribution in fragrant rice genotypes and related consequences on yield and grain quality traits. *Journal of Chemistry* 2017:1405878. doi:10.1155/2017/1405878.
- Li, K., Cao, C., Ma, Y., Su, D., Li, J. 2019. Identification of cadmium bioaccumulation in rice (*Oryza sativa* L.) by the soil-plant transfer model and species sensitivity distribution. *Science of the Total Environment* 692:1022-1028. doi:10.1016/j.scitotenv.2019.07.091.
- Li, G.Z., Chen, S.J., Li, N.Y., Wang, Y.Y., Kang, G.Z. 2021. Exogenous glutathione alleviates cadmium toxicity in wheat by influencing the absorption and translocation of cadmium. *Bulletin of Environmental Contamination and Toxicology* 107:320-326. doi:10.1007/s00128-021-03283-8.
- Li, H., Luo, N., Li, Y.W., Cai, Q.Y., Li, H.Y., Mo, C.H., et al. 2017. Cadmium in rice: Transport mechanisms, influencing factors, and minimizing measures. *Environmental Pollution* 224:622-630. doi:10.1016/j.envpol.2017.01.087.
- Liu, L.Z., Gong, Z.Q., Zhang, Y.L., Li, P.J. 2014. Growth, cadmium uptake and accumulation of maize *Zea mays* L. under the effects of arbuscular mycorrhizal fungi. *Ecotoxicology* 23:1979-1986. doi:10.1007/s10646-014-1331-6.
- Liu, L., Li, J.W., Yue, F.X., Yan, X.W., Wang, F.Y., Bloszies, S., et al. 2018. Effects of arbuscular mycorrhizal inoculation and biochar amendment on maize growth, cadmium uptake and soil cadmium speciation in Cd-contaminated soil. *Chemosphere* 194:495-503. doi:10.1016/j.chemosphere.2017.12.025.
- Liu, J., Shu, A., Song, W., Shi, W., Li, M., Zhang, W., et al. 2021. Long-term organic fertilizer substitution increases rice yield by improving soil properties and regulating soil bacteria. *Geoderma* 404:115287. doi:10.1016/j.geoderma.2021.115287.
- Liu, M., Sun, J., Li, Y., Xiao, Y. 2017. Nitrogen fertilizer enhances growth and nutrient uptake of *Medicago sativa* inoculated with *Glomus tortuosum* grown in Cd-contaminated acidic soil. *Chemosphere* 167:204-211. doi:10.1016/j.chemosphere.2016.09.145.
- MINAGRI. 2017. Boletín – Informe del arroz. Dirección General de Políticas Agrarias, Dirección de Estudios Económicos e Información Agraria. Ministerio de Agricultura y Riego (MINAGRI), Lima, Perú.
- Neira, E., Ramos, L., Razuri, L.R. 2020. Coeficiente del cultivo (Kc) del arroz a partir de lisímetro de drenaje en La Molina, Lima-Perú. *Idesia (Arica)* 38(2):49-55. doi:10.4067/S0718-34292020000200049.
- Newman, E.I. 1966. A method of estimating the total length of root in a sample. *Journal of Applied Ecology* 3(1):139-145. <https://www.jstor.org/stable/2401670>.
- Olsen, S.R., Sommers, L.E. 1982. Phosphorus. p. 403-430. In Page, A.L., et al. (eds.) *Methods of soil analysis: Part 2. Chemical and microbiological properties. Agronomy Monographs* 9. 2nd ed. ASA and SSSA, Madison, Wisconsin, USA. doi:10.2134/agronmonogr9.2.2ed.c24.

- Pandian, K., Gnasekaran, P., Subramaniayan, P., Chitraputhirapillai, S. 2016. Effect of biochar amendment on soil physical, chemical and biological properties and groundnut yield in rainfed Alfisol of semiarid tropics. *Archives of Agronomy and Soil Science* 62(9):1293-1310. doi:10.1080/03650340.2016.1139086.
- Phillips, J.M., Hayman, D.S. 1970. Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Transactions of the British Mycological Society* 55(1):158-161. doi:10.1016/s0007-1536(70)80110-3.
- Saikat, M., Arka, J., Chakraborty, A., Montakim, T., Talha, B., Firzan, N., et al. 2022. Impact of heavy metals on the environment and human health: Novel therapeutic insights to counter the toxicity. *Journal of King Saud University-Science* 34(3):101865. doi:10.1016/j.jksus.2022.101865.
- Song, W.S., Chen, S.B., Liu, J.F., Song, L., Li, N., Liu, B. 2015. Variation of Cd concentration in various rice cultivars and derivation of cadmium toxicity thresholds for paddy soil by species-sensitivity distribution. *Journal of Integrative Agriculture* 14(9):1845-1854. doi:10.1016/S2095-3119(14)60926-6.
- Song, Y., Wang, Y., Mao, W., Sui, H., Yong, L., Yang, D., et al. 2017. Dietary cadmium exposure assessment among the Chinese population. *PLOS ONE* 12:e0177978. doi:10.1371/journal.pone.0177978.
- Vaculik, M., Pavlovič, A., Lux, A. 2015. Silicon alleviates cadmium toxicity by enhanced photosynthetic rate and modified bundle sheath's cell chloroplasts ultrastructure in maize, *Ecotoxicology and Environmental Safety* 120:66-73. doi:10.1016/j.ecoenv.2015.05.026.
- Vallejos-Torres, G., Ruíz-Valles, R., Chappa-Santa María, C.E., Gaona-Jiménez, N., Marín, C. 2022. High genetic diversity in arbuscular mycorrhizal fungi influence cadmium uptake and growth of cocoa plants. *Bioagro* 34(1):75-84. doi:10.51372/bioagro341.7.
- Wang, Q.Q., Huang, Q., Guo, G.M., Qin, J.M., Luo, J.Y., Zhu, Z.Q., et al. 2020. Reducing bioavailability of heavy metals in contaminated soil and uptake by maize using organic-inorganic mixed fertilizer. *Chemosphere* 261:128122. doi:10.1016/j.chemosphere.2020.128122.
- Wu, Z.P., Wu, W.D., Zhou, S.L., Wu, S.H. 2016. Mycorrhizal inoculation affects Pb and Cd accumulation and translocation in pakchoi (*Brassica chinensis* L.) *Pedosphere* 26:13-26. doi:10.1016/S1002-0160(15)60018-2.
- Zhang, F.G., Liu, M.H., Li, Y., Che, Y.Y., Xiao, Y. 2019. Effects of arbuscular mycorrhizal fungi, biochar and cadmium on the yield and element uptake of *Medicago sativa*. *Science of the Total Environment* 655:1150-1158. doi:10.1016/j.scitotenv.2018.11.317.
- Zhang, J., Su, L., Yan, K., Li, M., He, Y., Zu, Y., et al. 2020. An arbuscular mycorrhizal fungus increased the macroaggregate proportion and reduced cadmium leaching from polluted soil. *International Journal of Phytoremediation* 23(7):684-692. doi:10.1080/15226514.2020.1849014.