

Phosphate solubilizing bacteria: Assessment of their effect on growth promotion and phosphorous uptake of mung bean (*Vigna radiata* [L.] R. Wilczek)

Buddhi Charana Walpola¹, and Min-Ho Yoon^{1*}

The aim of this work was to isolate phosphate solubilizing bacteria (PSB) and assess their effect on the growth of mung bean (*Vigna radiata* [L.] R. Wilczek) plants. Of 31 isolated PSB strains, two efficient strains, identified as *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2), were employed in further studies. Maximum P solubilization ($720.75 \mu\text{g mL}^{-1}$) was recorded from the cultures co-inoculated with *P. agglomerans* and *B. anthina*. A strong positive correlation was found between pH and soluble P concentration in the medium, as well as between titratable acidity and P solubilization. Both strains under greenhouse conditions remarkably enhanced shoot and root length, shoot and root dry matter, and P uptake of mung bean plants. Growth was found to be further improved by adding tricalcium phosphate (TCP) with PSB inoculation. Co-inoculation of both PSB strains and adding TCP exhibited the highest growth performances and P uptake of mung bean plants; this implies that their applicability as a promising alternative to minimize the P problem in agricultural soils.

Key words: *Burkholderia anthina*, co-inoculation, *Pantoea agglomerans*, phosphate solubilization, tricalcium phosphate, *Vigna radiata*.

INTRODUCTION

Despite the high total soil P content, plant P availability is often reported to be limited, particularly in tropical soils (Collavino et al., 2010). Most soil P is usually present as insoluble metal chelates (Vassilev et al., 2006); moreover, substantial amounts of applied chemical phosphate fertilizers are also rapidly converted into insoluble phosphate sources. This leads to regularly applying P fertilizers, which are not only costly, but also environmentally undesirable. In this context, microbial solubilization of soil insoluble phosphates into soluble forms is considered an important process in natural and agricultural ecosystems.

Several bacterial and fungal species with varied potentials to solubilize inorganic phosphates, also known as phosphate solubilizing microorganisms, have been found in the rhizosphere of plants. However, their number is not high enough to compete with other microbial species in the rhizosphere (Jain et al., 2012). Screening of potential phosphate solubilizing isolates, which can be used as bio-inoculants to increase plant growth and yield, is recognized as an area of interest because such microbial inoculants could substantially reduce the chemical fertilizer requirement. There are several previous reports

dealing with the application of phosphate solubilizing bacteria (PSB), either individually or combined, to assess their effects on the growth and biomass production of several crops (Fernández et al., 2007; Mittal et al., 2008; Vikram and Hamzehzarghani, 2008; Hariprasad and Niranjana, 2009; Jain et al., 2010). In the present study, we isolated and identified PSB strains and assessed their effect on the growth and nutrient uptake of mung bean plants cultivated under greenhouse conditions.

MATERIALS AND METHODS

Isolation and identification of bacterial strains

Soils employed to isolate bacterial strains were collected from Chungchugnam-do Province, Gongju-Gun area in South Korea. Moist field soil was mixed with a sterile 0.85% NaCl solution and shaken for 30 min. Serial dilutions were inoculated with NBRIP (National Botanical Research Institute Phosphorus) agar plates containing 10 g glucose, 5 g $\text{Ca}_3(\text{PO}_4)_2$, 5 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 0.25 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g KCl, and 0.1 g $(\text{NH}_4)_2\text{SO}_4$ in 1 L distilled water (Nautiyal, 1999). Medium pH was adjusted to 7 with HCl. Plates were incubated for 5 d at 30 °C. Colonies with clear halos were considered as phosphate solubilizing colonies (Vyas et al., 2007). A total of 31 bacterial isolates that exhibited clear zones on the agar plates were selected as phosphate solubilizing organisms. Based on the size of the clear zone, two bacterial isolates were selected as the efficient phosphate solubilizing organisms for further studies (PSB-1 and PSB-2).

Partial sequencing of 16S rRNA for the bacterial strains

¹Chungnam National University, College of Agriculture and Life Sciences, Daejeon, 305-764, Korea.

*Corresponding author (mhyoon@cnu.ac.kr).

Received: 3 January 2013.

Accepted: 22 July 2013.

doi:10.4067/S0718-58392013000300010

was performed with the help of a DNA sequencing service (Solgent, Daejeon, South Korea) with universal primers, 27F (5'-AGAGTTTGATCCTGGCTCAG-3'), and 1492R (5'-GGTTACCTTGTTACGACTT-3'). Polymerase chain reaction (PCR) was performed with initial denaturation at 95 °C for 2 min followed by 30 cycles with denaturation for 30 s at 94 °C, annealing for 30 s at 58 °C, and extension for 45 s at 72 °C. The final extension was held for 5 min at 72 °C. The online program BLAST (NCBI, 2012) was used in identifying the related sequences with known taxonomic information available at the databank of National Center for Biotechnology Information (NCBI, Bethesda, Maryland, USA). A phylogenetic tree was constructed with the CLUSTAL X program (Thompson et al., 1997), which involved sequence alignment by the neighbor-joining method (Saitou and Nei, 1987) and maximum parsimony with the MEGA4 program (Kumar et al., 2001). The grouping of sequences was based on confidence values obtained by bootstrap analysis of 1000 replicates. Gaps were edited in the BioEdit program (Hall, 1999) and evolutionary distances were calculated by Kimura's two-parameter model (Kimura, 1980). Reference sequences were retrieved from GenBank under the accession numbers indicated in the trees. The obtained sequences were deposited in the NCBI Genebank under accession numbers KF425001 (*Pantoea agglomerans*) and KF425002 (*Burkholderia anthina*).

Phosphate solubilization under *in vitro* conditions

Inorganic phosphate solubilization was assayed *in vitro* as single inoculation and co-inoculation. Both *P. agglomerans* and *B. anthina* strains were grown in sterilized liquid NBRIP medium (20 mL) at 30 °C for 2 d with continuous shaking at 150 rpm. Aliquots (1 mL) of each culture (10^8 CFU mL⁻¹ of each inoculant) were then transferred to a 500 mL flask (n = 3 per strain) containing sterilized liquid NBRIP medium (200 mL) and incubated for 3 d with continuous shaking at 30 °C. For co-inoculation, equal volumes (10^8 CFU mL⁻¹ of each inoculant) of *P. agglomerans* and *B. anthina* cultures were transferred to a 500 mL flask. Sterilized uninoculated medium served as a control. A sample (10 mL) of each culture and control were taken and centrifuged at 8000 rpm for 10 min. The clear supernatant was employed to determine the amount of P released into the medium. The culture medium pH was also recorded with the pH meter equipped with a glass electrode. Phosphorus availability was measured colorimetrically by the Murphy and Riley (1962) method. Bacteria growth was estimated by measuring absorbance at 660 nm. Samples from cultures grown in NBRIP medium were diluted 1:1 (v/v) with 1 N HCl to dissolve the residual insoluble phosphate and measured against a blank (Rodríguez et al., 2000). Titratable acidity produced by the strains during phosphate solubilization was determined by titration with 0.1 M NaOH (Takao, 1965).

Inoculum preparation for pot experiment

Single colonies of each strain were transferred to 500 mL flasks containing nutrient broth; colonies were then grown aerobically in flasks on a rotating shaker (150 rpm) for 48 h at 30 °C. The bacterial suspension was then diluted in sterile distilled water to a final concentration of 10^8 CFU mL⁻¹ and resulting suspensions were used to treat mung bean seeds. Surface sterilized seeds were soaked in separate bacterial suspensions approximately 30 min prior to planting. For dual inoculation, an equal volume (10^8 CFU mL⁻¹ of each inoculant) of two cultures were mixed and then employed to treat mung bean seeds (the same as for single inoculation).

Pot experiment

The experiment was carried out in a greenhouse located at Chungnam National University, South Korea. The potting soil was classified as sandy loam and had the following characteristics: pH 6.55, 300 mg NH₄⁺-N kg⁻¹, 300 mg NO₃⁻-N kg⁻¹, 255 mg P₂O₅ kg⁻¹ (Olsen P), and cation exchange capacity (CEC) of 10 cmol₍₊₎ L⁻¹. Pots (25 cm diameter, 35 cm height) were filled with this soil and basal N rates (50 mg kg⁻¹ soil) and K (120 mg kg⁻¹ soil) were applied as urea and potassium chlorite, respectively. Tricalcium phosphate (TCP) was supplied as soil P fertilizer at the rate of 160 mg kg⁻¹ based on the nutrient requirements of mung bean plants. Pots were arranged in a randomized complete block design with three replicates per treatment. The experimental plan was based on eight treatments as follows: (1) Control; (2) Soil + TCP; (3) Soil + PSB-1; (4) Soil + PSB-1 + TCP; (5) Soil + PSB-2; (6) Soil + PSB-2 + TCP; (7) Soil + PSB-1 + PSB-2; and (8) Soil + PSB-1 + PSB-2 + TCP.

Mung bean (*Vigna radiata* (L.) R. Wilczek var. Paiyur 1) seeds were surface-sterilized by immersing them in 0.1% sodium hypochlorite solution for 10 min and then washing them three times with distilled water. A 15 mm depth of soil was removed from the earthen pots and six seeds were placed at equal distances. Previously prepared 1 mL samples of each inoculant were uniformly applied on seeds as single and co-inoculation; seeds were then covered with a uniform 15 mm thick layer of soil. Control plants received 1 mL of diluted nutrient solution with no bacteria. Pots were watered daily to maintain soil field capacity during the study period. After 1 wk of germination, plants were thinned out and three plants per plot were left. Effects of promoting bacterial growth treatments were assessed by measuring main shoot and root length, shoot and root weight, and P uptake of mung bean plants 8 wk after planting. Root and shoot portions of plants were separated and air-dried before being kept in an oven at 70 °C to constant weight. Shoot and root dry weights were recorded separately and the mean weight of three plants was expressed as g plant⁻¹. Plant samples were finely ground after drying and employed to determine

plant P content by the Vanadate-Molybdate method described by Jackson (1973).

Analysis of pH, available P content, and PSB population in soil

The samples of rhizosphere soil were aseptically separated by shaking and washing roots with sterile distilled water to test soil pH, soil P content, and PSB population densities. Soil pH was measured in a 1:2.5 soil:water suspension with a pH meter (Model 440, Corning, Tewksbury Massachusetts, USA). Available soil P was extracted with sodium bicarbonate (pH 8.5) according to Olsen et al. (1954) and measured colorimetrically by the Murphy and Riley (1962) method.

The PSB population density was assessed by the pour plate method. The rhizosphere soil was collected by uprooting the plants. Soil adhering to the roots was serially diluted (10^2 to 10^6) with 0.85% NaCl solution and aliquots from 0.1 mL of the sample from each of these dilutions were spread on a petri dish containing NBRIP medium. Plates were incubated 3 d in an incubator at 30 °C. Colonies with clear halos were counted at the end of the incubation period.

Statistical analysis

Values were given as means \pm SD for triplicate samples. Data were subjected to ANOVA with the SAS package (SAS Institute, 1999). Duncan's Multiple Range Test (DMRT) was applied to test the significance of treatment means at $P \leq 0.05$.

RESULTS

Isolation and identification of bacterial strains

A total of 31 bacterial isolates that exhibited clear zones

on the agar plates were selected as phosphate solubilizing organisms. Two of these bacterial isolates were selected as efficient phosphate solubilizing organisms (PSB-1 and PSB-2). The two selected had a marked insoluble phosphate solubilizing ability as visualized by the clear zone development around the colonies after 3 d of incubation. According to 16S rRNA sequence analysis, strains were identified as *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2). Comparison of the 16S rRNA sequence among available strains of *Pantoea* and *Burkholderia* species showed high homology (> 99%) to *P. agglomerans* DSM3493 and *B. anthina* R4183. The neighbor-joining method was employed to construct the phylogenetic tree which illustrates the relationships of 16S rRNA strain sequence and other *Pantoea* and *Burkholderia* species (Figure 1).

Phosphate solubilization under *in vitro* conditions

Changes in NBRIP medium pH, microbial growth, titratable acidity, and soluble P content, released from inorganic P, due to the addition of single and co-inoculation of bacterial inoculants are shown in Table 1. Significant ($P \leq 0.05$) increments in soluble P content, titratable acid production, and microbial growth were observed with PSB inoculation. A significant reduction in the pH of the PSB-inoculated medium was also observed as compared with the control where it remained constant (pH 7). Co-inoculation of two PSB strains showed the highest phosphate solubilization when compared with single inoculation. A strong negative correlation between phosphate solubilization and pH, as well as a strong positive correlation between phosphate solubilization and microbial growth can also be observed (Table 2). The higher amount of P observed in the co-inoculated medium




Figure 1. Phylogenetic tree based on 16S rDNA gene sequences showing the position of *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2) strains with regard to related species. The scale bar indicates 0.02 substitutions per nucleotide position.

Table 1. Effects of single and co-inoculation of phosphate solubilizing bacteria (PSB) on phosphate solubilization.

Treatment	Amounts of soluble P	Dissolved	pH of medium	Microbial growth (optical density at 660 nm)	Titratable acidity
	$\mu\text{g mL}^{-1}$	%			
PSB-1	575.16 \pm 15.28b	57.56	3.21 \pm 0.12b	0.610 \pm 0.041b	30.67 \pm 1.22b
PSB-2	384.28 \pm 11.21c	38.24	3.54 \pm 0.11a	0.407 \pm 0.034c	18.67 \pm 1.01c
PSB-1 + PSB-2	720.75 \pm 14.37c	72.07	3.04 \pm 0.09c	0.764 \pm 0.038a	45.00 \pm 2.65a

Values are given as means \pm SD for triplicate samples. Means followed by the same letter(s) in each column are not significantly different at $P \leq 0.05$.

Table 2. Correlation coefficients of different parameters studied in single and co-inoculation of phosphate solubilizing bacteria (PSB).

	Soluble P	pH of medium	Microbial growth	Titratable acidity
Soluble P		-0.94**	0.92**	0.88**
pH of medium			-0.99**	-0.95**
Microbial growth				0.98**
Titratable acidity				

**Highly significant.

can be attributed to higher microbial growth, which seems to be influenced by the synergistic action of the two PSB strains.

Growth and P uptake in mung bean plants

Increased shoot length, root length, and shoot and root dry weight of mung bean plants were recorded from the seedlings raised with the PSB-inoculated seeds (Table 3). The best growth performances (24.16 cm, 27.22 cm, 2.88, and 2.93 g plant⁻¹ for shoot length, root length, shoot dry weight, and root dry weight, respectively) were recorded from the plants co-inoculated with *P. agglomerans* and *B. anthina* and amended with TCP. Although adding TCP resulted in better growth performances, no significant ($P \leq 0.05$) differences in shoot length, root length, and shoot-root dry weight were observed among soil treatments with and without TCP (Table 3).

As shown in Table 4, P uptake of mung bean plants showed a trend similar to the growth parameters. An increase in shoot P uptake, root P uptake, and total P uptake was observed in plants inoculated with *P. agglomerans* or *B. anthina* or both. Moreover, adding TCP to the PSB-inoculated seeds significantly ($P \leq 0.05$) increased shoot and root P uptake. Co-inoculation of PSB strains with TCP further improved P uptake as compared with single inoculation with any of the PSB strains and TCP

combination. Maximum P uptake (177.46 and 78.65 mg plant⁻¹ for shoot and root, respectively) was recorded for co-inoculated plants with TCP. There was no significant difference ($P \leq 0.05$) between uninoculated seeds treated with and without TCP.

Changes in pH, available soil P, and PSB population

The effect of single and co-inoculation of PSB strains on soil pH, available P content, and total PSB population are shown in Table 5. A more significant decrease ($P \leq 0.05$) in soil pH was recorded from PSB-inoculated soils than uninoculated soils. However, no significant ($P \leq 0.05$) difference in soil pH was observed among single and co-inoculated soils. Furthermore, available P content in the rhizosphere soil inoculated either by single PSB or both strains were found to be significantly ($P \leq 0.05$) higher than for uninoculated soil. This was further improved by adding TCP. The highest available P content (201.25 mg kg⁻¹ soil) recorded from co-inoculation of PSB strains with TCP was two times higher than for uninoculated soil. A

Table 4. Effect of *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2) on mung bean plant P uptake.

Treatment	P content in shoots	P content in roots	Total P uptake
	mg plant ⁻¹		
Control	136.55 \pm 2.47b	47.86 \pm 0.57d	184.41 \pm 3.57b
Soil + TCP	136.71 \pm 1.57b	46.66 \pm 1.12d	183.36 \pm 4.11b
Soil + PSB-1	141.12 \pm 2.87b	50.78 \pm 1.28dc	191.91 \pm 3.89b
Soil + PSB-1 + TCP	173.71 \pm 2.11a	73.18 \pm 1.07b	246.89 \pm 4.18a
Soil + PSB-2	140.91 \pm 1.58b	54.23 \pm 1.57c	195.13 \pm 2.84b
Soil + PSB-2 + TCP	172.59 \pm 2.87a	77.61 \pm 1.34ba	250.19 \pm 3.57a
Soil + PSB-1 + PSB-2	140.07 \pm 3.47b	54.02 \pm 1.27c	194.09 \pm 4.58b
Soil + PSB-1 + PSB-2+TCP	177.46 \pm 2.74a	78.65 \pm 1.12a	256.11 \pm 5.24a

Values are given as means \pm SD for triplicate samples. Means followed by the same letter(s) in each column are not significantly different at $P \leq 0.05$.

PSB: phosphate solubilizing bacteria; TCP: tricalcium phosphate.

Table 3. Effect of *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2) on mung bean plant growth.

Treatment	Shoot length	Root length	Shoot dry matter	Root dry matter
	cm plant ⁻¹		g plant ⁻¹	
Control	19.86 \pm 0.98cd	20.32 \pm 0.97 c	2.01 \pm 0.14ed	1.41 \pm 0.24 d
Soil + TCP	19.35 \pm 1.12d	20.01 \pm 1.25c	1.89 \pm 0.27e	1.47 \pm 0.24d
Soil + PSB-1	20.16 \pm 1.05cd	21.11 \pm 0.97bc	2.13 \pm 0.21d	1.81 \pm 0.21dc
Soil + PSB-1 + TCP	21.49 \pm 1.25bcd	22.64 \pm 1.24bc	2.39 \pm 0.25c	2.21 \pm 0.11bc
Soil + PSB-2	20.22 \pm 0.89cd	21.11 \pm 1.04bc	2.31 \pm 0.11c	1.88 \pm 0.14dc
Soil + PSB-2 + TCP	21.62 \pm 1.37bc	22.23 \pm 0.98bc	2.66 \pm 0.15b	2.68 \pm 0.15ba
Soil + PSB-1 + PSB-2	22.61 \pm 1.31ba	24.55 \pm 1.01ba	2.63 \pm 0.24b	2.61 \pm 0.26ba
Soil + PSB-1 + PSB-2+TCP	24.16 \pm 1.04a	27.22 \pm 1.15a	2.88 \pm 0.14a	2.93 \pm 0.28a

Values are given as means \pm SD for triplicate samples. Means followed by the same letter(s) in each column are not significantly different at $P \leq 0.05$.

PSB: phosphate solubilizing bacteria; TCP: tricalcium phosphate.

Table 5. Effect of *Pantoea agglomerans* (PSB-1) and *Burkholderia anthina* (PSB-2) on soil pH, available P content, and population of phosphate solubilizing bacteria (PSB) in rhizosphere soil of mung bean plants.

Treatment	Soil pH	Soil available P mg kg ⁻¹	nr of PSB CFU g ⁻¹ soil
Control	6.52 ± 0.32a	106.91 ± 1.35e	1.17 × 10 ³ d
Soil + TCP	6.53 ± 0.21a	104.39 ± 1.22e	1.26 × 10 ³ d
Soil + PSB-1	6.31 ± 0.35b	169.81 ± 2.34cbd	4.61 × 10 ⁴ d
Soil + PSB-1 + TCP	6.31 ± 0.12b	177.35 ± 3.01cb	5.41 × 10 ⁴ c
Soil + PSB-2	6.21 ± 0.11b	159.75 ± 1.57d	5.31 × 10 ⁴ d
Soil + PSB-2 + TCP	6.23 ± 0.24b	164.78 ± 2.24cd	5.68 × 10 ⁴ c
Soil + PSB-1 + PSB-2	6.28 ± 0.31b	182.39 ± 3.24b	6.91 × 10 ⁴ b
Soil + PSB-1 + PSB-2+TCP	6.28 ± 0.28b	201.25 ± 2.27a	8.36 × 10 ⁴ a

Values are given as means ± SD for triplicate samples. Means followed by the same letter(s) in each column are not significantly different at $P \leq 0.05$.

PSB: phosphate solubilizing bacteria; TCP: tricalcium phosphate.

remarkable increase in the PSB population was observed in PSB-inoculated rhizosphere soil when compared with uninoculated soil. The highest PSB population (8.36×10^6 CFU g⁻¹ soil) recorded from co-inoculation of PSB strains with TCP was approximately three times higher than for uninoculated soil.

DISCUSSION

Phosphate solubilization potential has been attributed to the strains' ability to reduce pH of the surroundings, either by releasing organic acids or protons (Hariprasad and Niranjana, 2009). Organic acids, such as gluconic acid, oxalic acid, and citric acid, secreted by PSB can directly solubilize mineral phosphate as a result of anion exchange or indirectly chelate both Fe and Al ions associated with phosphate. This leads to increased P availability, which ultimately increases plant P uptake. Studies on agar plates revealed that phosphate solubilizing microorganisms formed clear zones by solubilizing suspended TCP due to the release of organic acids into the surrounding medium (Gaur, 1990). In the present study, 31 PSB strains were isolated and two of these efficient PSB strains (*P. agglomerans* PSB-1 and *B. anthina* PSB-2) were selected for further studies; they had a marked insoluble phosphate solubilizing ability as visualized by the clear zone developed around the colonies. Previous reports also described some *Burkholderia* and *Pantoea* strains as being efficient phosphate solubilizers (Peix et al., 2001; Caballero-Mellado et al., 2007; Torres et al., 2008; Viruel et al., 2011; Khalimi et al., 2012; Silini-Cherif et al., 2012).

Titrate acidity is the total H⁺ concentration which can be produced by different compounds (Whitelaw et al., 1999). Organic acid production due to PSB inoculation should be predictable from titrate acidity in the culture medium. From the results of the present study, it can be reaffirmed that *P. agglomerans* and *B. anthina* were involved in organic acid production. The negative correlation between pH and soluble P content of the medium, as well as the positive correlation between soluble P content and titrate acid production, suggested

that acidification of the medium can facilitate phosphate solubilization. Co-inoculation showed a higher phosphate solubilizing ability than single inoculation; this suggests that both strains acted synergistically in phosphate solubilization. Yu et al. (2011) also reported similar findings after inoculating *Pseudomonas chlororaphis* and *Bacillus megaterium*.

Increased growth and P uptake of several crop plants due to PSB inoculation have been reported in a number of studies conducted under both growth chamber and greenhouse conditions (Dey et al., 2004; Fernández et al., 2007; Vikram and Hamzehzarghani, 2008; Hariprasad and Niranjana, 2009; Yu et al., 2011). The increase in shoot length, root length, shoot dry weight, and root dry weight of mung bean plants inoculated with PSB strains could be attributed to a greater absorption of nutrients, especially P. Compared with single inoculation, co-inoculation showed higher growth performances and P uptake; this suggests that both strains acted synergistically with each other to promote mung bean plant growth. However, phosphate solubilization is not the only way of promoting plant growth by PSB because they help plant growth by stimulating the efficiency of plant hormone production, such as auxins, cytokinins, gibberellins, and also some volatile compounds (Podile and Kishore, 2006). Enhanced plant growth after inoculation of PSB strains can be attributed to the ability of the strains to make P available and to simultaneously produce plant growth-promoting substances (Khalid et al., 2004; Linu et al., 2009; Ali et al., 2010).

Both strains used in this study exhibited the capacity to produce indoleacetic acid (data not shown); therefore, it might have contributed to enhanced shoot and root length through cell elongation and multiplication. A similar increase in growth and P uptake of mung bean plants due to inoculation of PSB strains was observed by Singh and Kapoor (1999), Vikram and Hamzehzarghani (2008), Ghanem and Abbas (2009), and Jha et al. (2011). Ghanem and Abbas (2009) observed an increase in plant height, number of branches, number of pods, grain weight, and eventually higher seed and straw yields in mung bean plants after inoculation of *B. megaterium* in salt-affected soils. Increased growth and P uptake have been reported for *Azotobacter chroococcum* in wheat (Kumar et al., 2001), *Pseudomonas fluorescens* in peanut (Dey et al., 2004), *Pseudomonas* species in wheat (Babana and Antoun, 2006), *Pseudomonas* species and *Bacillus cereus* in walnut (Yu et al., 2011), and *Paenibacillus polymyxa* and *B. megaterium* in tomato (El-Yazeid and Abou-Aly, 2011). According to Fernández et al. (2007), shoot length of soybean plants increased after inoculation of *Burkholderia* sp. PER2F by 40% and 60% when compared with uninoculated soil/seed and uninoculated soil/seed treated with soluble P, respectively. However, an increase in height and biomass, but not P content, of canola plants was observed by de Freitas et al. (1997) after inoculation

of *Bacillus* and *Xanthomonas*, which are two PSB strains.

Results of the present study as to maximum plant growth and P uptake recorded with co-inoculation of two PSB strains with TCP are in line with the findings of Qureshi et al. (2011), who also observed similar results when co-inoculating phosphate solubilizing and nodule-forming bacteria *Rhizobium phaseoli* and *B. megaterium* in mung bean plants. As for their observations, single rhizobium inoculation resulted in pod and straw yield of 24.0 and 30.20 g pot⁻¹, respectively, whereas the respective figures were 24.3 and 32.07 g pot⁻¹ for co-inoculation. Co-inoculation also produced higher root mass (231.3 g), root length (50.54 cm), nodule number (78), and nodular mass (0.216 g) as compared with the control.

Adding an insoluble phosphate source significantly increased total PSB populations in the soil, which implies that adding TCP was obviously beneficial for these isolates to proliferate and survive. Hence, more available P would be released into the soil and utilized by mung bean plants. A similar increase in the PSB population and available P content was observed by Yu et al. (2011). They also observed a positive correlation between available soil P content and PSB populations. A positive and significant correlation between phosphate solubilizing microorganisms and soil organic matter content was previously reported by Venkateswarlu et al. (1984). Other researchers (Vyas and Gulati, 2009) also reported decreased soil pH in PSB-inoculated soil and indicated organic acid production of both strains. Soil pH reduction was found to be much lower than in the culture medium (Table 1), which could be due to the buffering nature of the soil used in the experiment (Gyaneshwar et al., 1998).

CONCLUSIONS

This study has provided ample evidence to prove the strains' capacity to enhance plant growth. Co-inoculation of two PSB strains acted synergistically with each other and this was responsible for the increase in several growth parameters as compared with single inoculation. Nonetheless, further studies are needed under field conditions to confirm the present findings and their eventual commercial applications.

LITERATURE CITED

- Ali, B., A.N. Sabri, and S. Hasnain. 2010. Rhizobacterial potential to alter auxin content and growth of *Vigna radiata* (L.) World Journal of Microbiology and Biotechnology 26:1379-1384.
- Babana, A.H., and H. Antoun. 2006. Biological system for improving the availability of Tilemsi phosphate rock for wheat (*Triticum aestivum* L.) cultivated in Mali. Nutrient Cycling in Agroecosystems 76:285-295.
- Caballero-Mellado, J., J. Onofre-Lemus, P. Estrada-de Los Santos, and L. Martínez-Aguilar. 2007. The tomato rhizosphere, an environment rich in nitrogen-fixing *Burkholderia* species with capabilities of interest for agriculture and bioremediation. Applied and Environmental Microbiology 73:5308-5319.
- Collavino, M.M., P.A. Sansberro, L.A. Mroginski, and O.M. Aguilar. 2010. Comparison of in vitro solubilization activity of diverse phosphate-solubilizing bacteria native to acid soil and their ability to promote *Phaseolus vulgaris* growth. Biology and Fertility of Soils 46:727-738.
- de Freitas, J.R., M.R. Banerjee, and J.J. Germida. 1997. Phosphate-solubilizing rhizobacteria enhance the growth and yield but not phosphorus uptake of canola (*Brassica napus* L.) Biology and Fertility of Soils 24:358-364.
- Dey, R., K.K. Pal, D.M. Bhatt, and S.M. Chauhan. 2004. Growth promotion and yield enhancement of peanut (*Arachis hypogaea* L.) by application of plant growth-promoting rhizobacteria. Microbiological Research 159:371-394.
- El-Yazeid, A.A., and H.E. Abou-Aly. 2011. Enhancing growth, productivity and quality of tomato plants using phosphate solubilizing microorganisms. Australian Journal of Basic and Applied Sciences 7:371-379.
- Fernández, L.A., P. Zalba, M.A. Gómez, and M.A. Sagardoy. 2007. Phosphate-solubilization activity of bacterial strains in soil and their effect on soybean growth under greenhouse conditions. Biology and Fertility of Soils 43:805-809.
- Gaur, A.C. 1990. Phosphate solubilizing microorganisms as biofertilizers. Omega Scientific Publishers, New Delhi, India.
- Ghanem, K.H.M., and E.E. Abbas. 2009. Improvement of mung bean growth and productivity in salinity-affected soil after seed inoculation with phosphate-dissolving bacteria. African Crop Science Conference Proceedings 9:385-389.
- Gyaneshwar, P., G. Naresh Kumar, and L.J. Parekh. 1998. Effect of buffering on the phosphate-solubilizing ability of microorganisms. World Journal of Microbiology and Biotechnology 14:669-673.
- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41:95-98.
- Hariprasad, P., and S.R. Niranjana. 2009. Isolation and characterization of phosphate solubilizing rhizobacteria to improve plant health of tomato. Plant and Soil 316:13-24.
- Jackson, M.L. 1973. Soil chemical analysis. Prentice-Hall, New Delhi, India.
- Jain, R., J. Saxena, and V. Sharma. 2010. The evaluation of free and encapsulated *Aspergillus awamori* for phosphate solubilization in fermentation and soil-plant system. Applied Soil Ecology 46:90-94.
- Jain, R., J. Saxena, and V. Sharma. 2012. Effect of phosphate-solubilizing fungi *Aspergillus awamori* S29 on mungbean (*Vigna radiata* cv. RMG 492) growth. Folia Microbiologica 57:533-541.
- Jha, A., D. Sharma, and J. Saxena. 2011. Effect of single and dual phosphate solubilizing bacterial strain inoculations on overall growth of mung bean plants. Archives of Agronomy and Soil Science 58:967-981.
- Khalid, A., M. Arshad, and Z.A. Zahir. 2004. Screening plant growth promoting rhizobacteria for improving growth and yield of wheat. Journal of Applied Microbiology 96:473-480.
- Khalimi, K., D.N. Suprapt, and Y. Nitta. 2012. Effect of *Pantoea agglomerans* on growth promotion and yield of rice. Agricultural Science Research Journals 2:240-249.
- Kimura, M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. Journal of Molecular Evolution 16:111-120.
- Kumar, V., R.K. Behl, and N. Narula. 2001. Establishment of phosphate-solubilizing strains of *Azotobacter chroococcum* in the rhizosphere and their effect on wheat cultivars under green house conditions. Microbiological Research 156:87-93.
- Kumar, S., K. Tamura, I.B. Jakobsen, and M. Nei. 2001. MEGA2: molecular evolutionary genetics analysis software. Bioinformatics 17:1244-1245.
- Linu, M.S., J. Stephen, and M.S. Jisha. 2009. Phosphate solubilizing *Gluconacetobacter* sp., *Burkholderia* sp. and their potential interaction with cowpea (*Vigna unguiculata* (L.) Walp.) International Journal of Agricultural Research 4:79-87.

- Mittal, V., O. Singh, H. Nayyar, J. Kaur, and R. Tewari. 2008. Stimulatory effect of phosphate-solubilizing fungal strains (*Aspergillus awamori* and *Penicillium citrinum*) on the yield of chickpea (*Cicer arietinum* L. cv. GPF2). *Soil Biology and Biochemistry* 40:718-727.
- Murphy, J., and J.P. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta* 27:31-36.
- Nautiyal, C.S. 1999. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiology Letters* 170:265-270.
- NCBI. 2012. BLAST: Basic Local Alignment Search Tool. National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA. Available at <http://www.ncbi.nlm.nih.gov/BLAST> (accessed on 28 November 2012).
- Olsen, S.R., C.V. Cole, F.S. Watanabe, and L.A. Dean. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939. 19 p. United States Department of Agriculture (USDA), Washington, DC., USA.
- Peix, A., P.F. Mateos, C. Rodríguez-Barrueco, E. Martínez-Molina, and E. Velazquez. 2001. Growth promotion of common bean (*Phaseolus vulgaris* L.) by a strain of *Burkholderia cepacia* under growth chamber conditions. *Soil Biology and Biochemistry* 33:1927-1935.
- Podile, A.R., and G.K. Kishore. 2006. Plant growth promoting rhizobacteria (PGPR). p. 195-230. In Gnanamanickam, S.S (ed.) *Plant-associated bacteria*. Springer, Dordrecht, The Netherlands.
- Qureshi, M.A., M.A. Shakir, A. Iqbal, N. Akhtar, and A. Khan. 2011. Co-inoculation of phosphate solubilizing bacteria and rhizobia for improving growth and yield of mungbean (*Vigna radiata* L.) *Journal of Animal and Plant Sciences* 21:491-497.
- Rodríguez, H., T. Gonzalez, and G. Selman. 2000. Expression of a mineral phosphate solubilizing gene from *Erwinia herbicola* in two rhizobacterial strains. *Journal of Biotechnology* 84:155-161.
- Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 4:406-425.
- SAS Institute. 1999. SAS/STAT User's guide version 8. SAS Institute, Cary, North Carolina, USA.
- Silini-Cherif, H., A. Silini, M. Ghoul, and S. Yadav. 2012. Isolation and characterization of plant growth promoting traits of a rhizobacteria: *Pantoea agglomerans* Ima2. *Pakistan Journal of Biological Sciences* 15:267-276.
- Singh, S., and K.K. Kapoor. 1999. Inoculation with phosphate-solubilizing microorganisms and a vesicular-arbuscular mycorrhizal fungus improves dry matter yield and nutrient uptake by wheat grown in a sandy soil. *Biology and Fertility of Soils* 28:139-144.
- Takao, S. 1965. Organic acid production by basidiomycetes. I. Screening of acid-producing strains. *Applied Microbiology* 13:732-737.
- Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougin, and D.G. Higgins. 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25:4876-4882.
- Torres, A.R., W.L. Araújo, L. Cursino, M. Hungria, F. Plotegher, F.L. Mostasso, and J.L. Azevedo. 2008. Diversity of endophytic enterobacteria associated with different host plants. *Journal of Microbiology* 46:373-379.
- Vassilev, N., M. Vassileva, and I. Nikolaeva. 2006. Simultaneous P-solubilizing and biocontrol activity of microorganisms: potentials and future trends. *Applied Microbiology and Biotechnology* 71:137-144.
- Venkateswarlu, B., A.V. Rao, and O. Raina. 1984. Evaluation of phosphorus solubilisation by microorganisms isolated from Aridisols. *Journal of the Indian Society of Soil Science* 32:273-277.
- Vikram, A., and H. Hamzehzarghani. 2008. Effect of phosphate solubilizing bacteria on nodulation and growth parameters of greengram (*Vigna radiata* L. Wilczek). *Research Journal of Microbiology* 3:62-72.
- Viruel, E., M.E. Lucca, and F. Siñeriz. 2011. Plant growth promotion traits of phosphobacteria isolated from Puna, Argentina. *Archives of Microbiology* 193:489-496.
- Vyas, P., and A. Gulati. 2009. Organic acid production in vitro and plant growth promotion in maize under controlled environment by phosphate-solubilizing fluorescent *Pseudomonas*. *BMC Microbiology* 9:174-189.
- Vyas, P., P. Rahi, A. Chauhan, and A. Gulati. 2007. Phosphate solubilization potential and stress tolerance of *Eupenicillium parvum* from tea soil. *Mycological Research* 111:931-938.
- Whitelaw, M.A., T.J. Harden, and K.R. Helyar. 1999. Phosphate solubilisation in solution culture by the soil fungus *Penicillium radicum*. *Soil Biology and Biochemistry* 31:655-665.
- Yu, X., X. Liu, T.H. Zhu, G.H. Liu, and C. Mao. 2011. Isolation and characterization of phosphate-solubilizing bacteria from walnut and their effect on growth and phosphorus mobilization. *Biology and Fertility of Soils* 47:437-446.