

Susceptibility of *Spodoptera frugiperda* and *Helicoverpa gelotopoeon* (Lepidoptera: Noctuidae) to the entomopathogenic nematode *Steinernema diaprepesi* (Rhabditida: Steinernematidae) under laboratory conditions

Milena G. Caccia^{1*}, Eleodoro Del Valle², Marcelo E. Doucet¹, and Paola Lax¹

Spodoptera frugiperda Smith and *Helicoverpa gelotopoeon* (Dyar) are important agricultural pests of several crops. The aim of the present work was to evaluate the susceptibility of larvae of both insects to an isolate of *Steinernema diaprepesi* Nguyen & Duncan under laboratory conditions, as well as the capacity of the nematode to multiply on these lepidoterans. Larvae (n = 15) were exposed to 0 (control), 50, and 100 infective juveniles (IJs) per Petri dish. Mortality was evaluated every 24 h during 6 d, and emerging IJs were counted. Mortality of *S. frugiperda* was 93% and 100% with 50 and 100 IJs dosage, and 87% and 93% in *H. gelotopoeon*, respectively. The production of IJs was significantly different between doses ($P \leq 0.05$) for *S. frugiperda* (11 329 with 50 IJs vs. 27 155 with 100 IJs) but not for *H. gelotopoeon* (19 830 vs. 26 361, respectively). This is the first study evaluating the susceptibility of these lepidoterans to *S. diaprepesi*. These results encourage the possibility of using this nematode for biological control of both pests.

Key words: Infective juveniles, biological control.

INTRODUCTION

The family Noctuidae is the most diverse group within Lepidoptera and includes the highest number of species of agricultural importance (Specht et al., 2004). The fall armyworm *Spodoptera frugiperda* Smith is a polyphagous insect that causes losses in several crops, such as peanut, sugarcane, cotton, soybean, alfalfa and particularly corn (Casmuz et al., 2010). Its wide distribution extends from Argentina and Chile to the southeast of Canada (Ashley et al., 1989). It is the most important lepidopteran pest of maize in northern Argentina and different countries of the Neotropical region (Casmuz et al., 2010).

The bollworm, *Helicoverpa gelotopoeon* (Dyar), damages mainly pods and seeds of soybean and other legumes. It also affects cotton, onion, and sunflower, among other crops (Specht et al., 2004). It is present in Argentina, Brazil, Chile, Uruguay, and Paraguay (Pastrana, 2004; Fichetti et al., 2009). In central-western Córdoba Province (Argentina), it can cause great damage to soybean seedlings.

The use of synthetic pesticides to control Lepidoptera worldwide has contributed to the development of

resistance to insecticides in several species (Bloem and Carpenter, 2001), and generated environmental pollution. Hence, alternative management strategies are currently being explored, such as: the use of viruses (Escribano et al., 1999), fungi (Lacey et al., 2001) and transgenic cultivars with toxins of *Bacillus thuringiensis* (Romeis et al., 2006). Also, entomopathogenic nematodes (EPNs) (Heterorhabditidae and Steinernematidae) are used against insect pests (Divya and Sankar, 2009). The infective juveniles (IJs) penetrate the insect hemocoel and release their symbiotic bacteria (*Xenorhabdus* spp. in Steinernematidae and *Photorhabdus* spp. in Heterorhabditidae), which multiply and generate metabolites that kill the insect and serve as source of food for nematodes (Dowds and Peters, 2002). Susceptibility of *S. frugiperda* to EPNs has been frequently reported (Molina Ochoa et al., 1996; Doucet et al., 1999; García et al., 2008; Andaló et al., 2010), but to present, no studies have evaluated the effect of *Steinernema diaprepesi* Nguyen & Duncan. Furthermore, infectivity of EPNs on *H. gelotopoeon* has still not been reported. The aim of the present work was to evaluate the susceptibility of *S. frugiperda* and *H. gelotopoeon* larvae to an *S. diaprepesi* isolate under laboratory conditions, as well as the capacity of the parasite to multiply on the pests.

MATERIALS AND METHODS

An *S. diaprepesi* isolate (strain SRC) obtained from soil samples from Santa Rosa de Calchines, Department of Garay (Santa Fe Province, Argentina) (Lax et al., 2011) was

¹Universidad Nacional de Córdoba, Facultad de Ciencias Exactas, Físicas y Naturales, Rondeau 798, X5000, Córdoba, Argentina.
*Corresponding author (milencaccia@yahoo.com.ar).

²Universidad Nacional del Litoral, Facultad de Ciencias Agrarias, 3080, Esperanza, Santa Fe, Argentina.

Received: 28 May 2013.

Accepted: 4 November 2013.

doi:10.4067/S0718-58392014000100019

used. Nematodes were maintained on larvae of *Galleria mellonella* L. (Lepidoptera: Pyralidae) in the laboratory, following the technique of Kaya and Stock (1997). The IJs were collected using White (1927) traps and kept in water at 10 ± 2 °C until use, for no longer than 21 d.


Same size last-instar of *S. frugiperda* and *H. gelotopoeon* larvae, provided by Laboratorio Líder (Córdoba Province, Argentina), were used. Infections were performed individually in Petri dishes (35 mm diameter) by applying IJs immersed in 0.5 mL water at 50 and 100 IJs per dish onto double filter paper. The same volume of water without nematodes was used as control. Each treatment contained 15 larvae; insects were maintained in the dark at 25 °C and mortality was evaluated every 24 h for 6 d. Four days after death, larvae were individually transferred to White traps and kept at 25 °C. Emerging IJs were collected the following 10 d and counted. For this, suspensions were increased to a volume of 10 mL and homogenized by bubbling; three 1-mL aliquots were taken from each suspension and placed for observation in a counting chamber under light microscope.

Insect accumulated mortality was compared between dosages in each day of observation and between species on day 6. For analysis, generalized linear models with binomial distribution and logit link function were performed with SPSS (2000). The number of IJs produced per larva was subjected to ANOVA ($P \leq 0.05$); differences between treatments were determined with a Fisher test (InfoStat, 2002).

RESULTS AND DISCUSSION

The isolate of *S. diaprepesi* used was able to infect and reproduce on both insect species. Larvae death started on day 2 in all treatments (Figure 1). At the end of the experiment, mortality of *S. frugiperda* was 93% at 50 IJ dosages, whereas the higher dosage reached 100% mortality on day 4. Mortality of *H. gelotopoeon* was 87% (50 IJs) and 93% (100 IJs). Control treatments had no mortality. No differences in mortality between insect species occurred in the 6 d analyzed or in accumulated mortality on day 6. The production of IJs was significantly different between dosages for *S. frugiperda*, but not for *H. gelotopoeon* (Table 1). No differences in IJ production occurred between insect species inoculated with the same EPN concentration.

This is the first study evaluating the susceptibility of *S. frugiperda* to *S. diaprepesi*. This insect has been previously found to be susceptible to other EPN species at different doses. In the laboratory, Andaló et al. (2010) studied the efficiency of 17 EPN populations on *S. frugiperda* at various concentrations; at the lowest dose (100 IJs), mortality reached 40-85%, with a population of *Heterorhabditis* sp. being the most efficient, not significant differences in mortality occurred with an increased dose of that EPN isolate. Molina Ochoa et al. (1996) analyzed



IJs: infective juveniles.

Figure 1. Accumulated mortality of *Spodoptera frugiperda* (A) and *Helicoverpa gelotopoeon* (B) larvae during 6 d post-inoculation with *Steinernema diaprepesi* strain SRC.

six EPN species against the same lepidopteran; at 100 IJs, the highest mortality (81%) occurred with an isolate of *Steinernema carpocapsae* (Filipjev), whereas an isolate of *Heterorhabditis bacteriophora* Poinar caused only 65% mortality.

Although there are no published studies on the effect of EPNs on *H. gelotopoeon*, other species of the genus have been studied. Mortality of *Helicoverpa armigera* (Hübner) larvae inoculated with 50 and 100 IJs of three EPN species ranged between 11% and 51% after 5 d (Kary et al., 2012). Five days post-inoculation (DPI) with IJs of *Steinernema riobravisi* Poinar & Raulston at 40 and 100 IJs, Cabanillas and Raulston (1994) observed that prepupae of *Helicoverpa zea* (Boddie) exhibited similar susceptibility to that obtained herein with *H. gelotopoeon* (85% and 100%, respectively).

It can be assumed that the final EPN population varies with the number of IJs inoculated (Koppenhöfer and Kaya, 1995). Herein, differences in final population between dosages were detected in *S. frugiperda* but not in *H. gelotopoeon*. Several unknown factors affect

Table 1. Final population of *Steinernema diaprepesi* strain SRC collected 10 d after death of *Spodoptera frugiperda* and *Helicoverpa gelotopoeon* larvae (n = 15) under two infective juvenile (IJ) dosages.

Treatments (IJ dosages)	<i>S. frugiperda</i>	<i>H. gelotopoeon</i>
50	11 329 ± 7473Aa	19 830 ± 27 691Aa
100	27 155 ± 8855Ba	26 361 ± 26 926Aa

Different uppercase letter indicates significant differences between rows (dosages) and the same lower-case letter indicates non significant differences between columns (species) according to Fisher's test ($P \leq 0.05$).

LITERATURE CITED

- nematode production *in vivo*, and these factors interact in unpredictable ways (Dolinski et al., 2007). This may help explain the high variability in the number of IJs produced per larva, especially in *H. gelotopoeon*. The amount of IJs obtained with both insects was lower than that observed in other studies with different hosts. On *Anticarsia gemmatilis* Hübner adults, Cagnolo et al. (2011) obtained 66 600 and 58 000 IJs of *Steinernema rarum* (Doucet) at 500 and 50 IJs, respectively, 10 d after insect death. As in our study, no significant differences in final IJ population occurred between dosages. At a higher dosage (200 IJs) of *Heterorhabditis baujardi* Phan, Subbotin, Nguyen & Moens on *G. mellonella*, Dolinski et al. (2007) obtained 165 000 IJs after recovering emerging nematodes for 7 d. Although these yields were higher than those observed herein, the importance of our results lie in that they confirm the capacity of the *S. diaprepesi* isolate to complete its life cycle successfully in both hosts evaluated.
- The IJ production of *S. riobravus* per *H. zea* larva peaked 375 000 with 40 IJs, with a reduction observed with higher dosages (Cabanillas and Raulston, 1994). Likewise, with *G. mellonella* larvae, the final population (32 000 IJs) of *Heterorhabditis megidis* Poinar, Jackson & Klein 5 d after the start of IJ emergence was obtained by inoculating 300 IJs, and decreased with an increase in dosage (Boff et al., 2000). This lower final population with an increased IJ inoculum might be attributed to a higher intraspecific competition among nematodes (Selvan et al., 1993). In our work, this trend was not observed; however, this should be considered when the optimal dose of inoculum is estimated for a host species.
- Besides, the effect of the isolate needs to be evaluated in the field, considering other variables, such as nematode capacity to find the host, influence of temperature and humidity (Rohde et al., 2010), and interaction with other pathogens (García et al., 2008). In addition, the site of the host plant where each stage of the insect life cycle occurs should be taken into account to develop an effective EPN application strategy.
- ### CONCLUSIONS
- Our results show that both lepidopterans, *Spodoptera frugiperda* and *Helicoverpa gelotopoeon*, were suitable hosts for *Steinernema diaprepesi* strain SRC. The nematode completed its life cycle and produced a high number of IJs, showing potential for biological control within an integrated pest management.
- ### ACKNOWLEDGEMENTS
- The authors thank the support of Secretaría de Ciencia y Técnica (Universidad Nacional de Córdoba), the Ministerio de Ciencia y Tecnología (Province of Córdoba), and the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).
- Andaló, V., V. Santos, G. Furtado Moreira, C. Costa Moreira, and J.A. Moino. 2010. Evaluation of entomopathogenic nematodes under laboratory and greenhouses conditions for the control of *Spodoptera frugiperda*. *Ciência Rural* 40:1860-1866.
- Ashley, T.R., B.R. Wiseman, F.M. Davis, and K.L. Andrews. 1989. The fall armyworm; a bibliography. *Florida Entomologist* 72:152-202.
- Bloem, S., and J.E. Carpenter. 2001. Population suppression by irradiated Lepidoptera. *Florida Entomologist* 84:165-171.
- Boff, M.I.C., G.L. Wiegler, L.J.M. Gerritsen, and P.H. Smits. 2000. Development of the entomopathogenic nematode *Heterorhabditis megidis* strain NLH-E 87.3 in *Galleria mellonella*. *Nematology* 2:303-308.
- Cabanillas, H.E., and J.R. Raulston. 1994. Pathogenicity of *Steinernema riobravus* against corn earworm, *Helicoverpa zea* (Boddie). *Fundamental and Applied Nematology* 17:219-223.
- Cagnolo, S.R., M.L. Peschiutta, and M.A. Bertolotti. 2011. Susceptibility of adults of *Anticarsia gemmatilis* Hübner, 1818 (Lepidoptera: Noctuidae) to the entomopathogenic nematode *Steinernema rarum* (Doucet, 1986) Mamiya, 1988 (Rhabditida: Steinernematidae) under laboratory conditions. *Nematology* 13:373-376.
- Casmuz, A., M.G. Juárez Socías, M.G. Murúa, S. Prieto, S. Medina, E. Willink, et al. 2010. Revisión de los hospederos del gusano cogollero del maíz, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Revista de la Sociedad Entomológica Argentina* 69:209-231.
- Divya, K., and M. Sankar. 2009. Entomopathogenic nematodes in pest management. *Indian Journal of Science and Technology* 2:53-60.
- Dolinski, C., E.E. Del Valle, R.S. Burla, and I. Ribeiro Machado. 2007. Biological traits of two native Brazilian entomopathogenic nematodes (Rhabditida: Heterorhabditidae). *Nematologia Brasileira* 31:180-185.
- Doucet, M.M.A. de, M.A. Bertolotti, A.L. Giayetto, and M.B. Miranda. 1999. Host range, specificity, and virulence of *Steinernema feltiae*, *Steinernema rarum*, and *Heterorhabditis bacteriophora* (Steinernematidae and Heterorhabditidae) from Argentina. *Journal of Invertebrate Pathology* 73:237-242.
- Dowds, B.C.A., and A. Peters. 2002. Virulence mechanisms. p. 79-98. In R. Gaugler (ed.) *Entomopathogenic Nematology*. CABI, New York, USA.
- Escribano, A., T. Williams, D. Goulson, R.D. Cave, J.W. Chapman, and P. Caballero. 1999. Biological and microbial control selection of a nucleopolyhedrovirus for control of *Spodoptera frugiperda* (Lepidoptera: Noctuidae): structural, genetic, and biological comparison of four isolates from the Americas. *Journal of Economic Entomology* 92:1079-1085.
- Fichetti, P., S. Avalos, V. Mazzuferi, and J. Carreras. 2009. Lepidoptera presentes en el cultivo de garbanzo en Córdoba, Argentina. *Boletín de Sanidad Vegetal. Plagas* 35:49-58.
- García, L.C., C.G. Raetano, and L.G. Leite. 2008. Application technology for the entomopathogenic nematodes *Heterorhabditis indica* and *Steinernema* sp. (Rhabditida: Heterorhabditidae and Steinernematidae) to control *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) in corn. *Neotropical Entomology* 37:305-311.
- InfoStat. 2002. InfoStat version 1.1. Grupo InfoStat, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, Argentina.
- Kary, N.E., A. Golizadeh, H.R. Dastjerdi, D. Mohammadi, S. Afghahi, M. Omrani, et al. 2012. A laboratory study of susceptibility of *Helicoverpa armigera* (Hübner) to three species of entomopathogenic nematodes. *Munis Entomology & Zoology* 7:372-379.
- Kaya, H., and P. Stock. 1997. Techniques in insect nematology. p. 281-324. In L.A. Lacey (ed.) *Biological techniques: Manual of techniques in insect pathology*. Academic Press, Millbrae, California, USA.

- Koppenhöfer, A.M., and H.K. Kaya. 1995. Density-dependent effects on *Steinernema glaseri* (Nematoda: Steinernematidae) within an insect host. *Journal of Parasitology* 81:797-799.
- Lacey, L.A., R. Frutos, H.K. Kaya, and P. Vail. 2001. Insect pathogens as biological control agents: do they have a future? *Biological Control* 21:230-248.
- Lax, P., E. del Valle, J. Rondan Dueñas, M. Caccia, C. Gardenal, and M.E. Doucet. 2011. First record of *Steinernema diaprepesi* (Steinernematidae) in Argentina. *Nematropica* 41:377.
- Molina Ochoa, J., J.J. Hamm, R. Lezama Gutierrez, L.F. Bojalil Jaber, M. Arenas Vargas, and M. González Ramírez. 1996. Virulence of six entomopathogenic nematodes (Steinernematidae and Heterorhabditidae) on immature stages of *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Vedalia Revista Internacional de Control Biológico* 3:25-29.
- Pastrana, J.A. 2004. Los lepidópteros argentinos: sus plantas hospedadoras y otros sustratos alimenticios. Sociedad Entomológica Argentina, Tucumán, Argentina.
- Rohde, C., A. Moino, M.A.T. da Silva, F.D. Carvalho, and C.S. Ferreira. 2010. Influence of soil temperature and moisture on the infectivity of entomopathogenic nematodes (Rhabditida: Heterorhabditidae, Steinernematidae) against larvae of *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae). *Neotropical Entomology* 39:608-611.
- Romeis, J., M. Meissle, and F. Bigler. 2006. Transgenic crops expressing *Bacillus thuringiensis* toxins and biological control. *Nature Biotechnology* 24:63-71.
- Selvan, S., J.F. Campbell, and R. Gaugler. 1993. Density-dependent effects on entomopathogenic nematodes (Heterorhabditidae and Steinernematidae) within an insect host. *Journal of Invertebrate Pathology* 62:278-284.
- Specht, A., E.J.E. Silva, e D. Link. 2004. Noctuídeos (Lepidoptera, Noctuidae) do museu entomológico Ceslau Biezanko, Departamento de Fitossanidade, Faculdade de Agronomia "Eliseu Maciel", Universidade Federal de Pelotas, Rio Grande do Sul. *Revista Brasileira de Agrociência* 10:389-409.
- SPSS. 2000. Version 10.0. Syntax reference guide. SPSS Inc., Chicago, Illinois, USA.
- White, G.F. 1927. A method for obtaining infective nematode larvae from cultures. *Science* 66:302-303.